

## ribG

### RELATED APPLICATIONS

- 5 This application claims benefit of PCT Application, Number PCT/US97/02318, filed February 19, 1997.

### FIELD OF THE INVENTION

- 10 This invention relates to newly identified polynucleotides and polypeptides, and their production and uses, as well as their variants, agonists and antagonists, and their uses. In particular, the invention relates to novel polynucleotides and polypeptides of the riboflavin specific deaminase family, hereinafter referred to as "ribG".

### BACKGROUND OF THE INVENTION

- 15 It is particularly preferred to employ Staphylococcal genes and gene products as targets for the development of antibiotics. The Staphylococci make up a medically important genera of microbes. They are known to produce two types of disease, invasive and toxigenic. Invasive infections are characterized generally by abscess formation effecting both skin surfaces and deep tissues. S. aureus is the second leading cause of bacteremia in cancer patients. Osteomyelitis, septic arthritis, septic thrombophlebitis and acute bacterial endocarditis are also relatively common. There are at least three clinical conditions resulting from the toxigenic properties of Staphylococci. The manifestation of these diseases result from the actions of exotoxins as opposed to tissue invasion and bacteremia. These conditions include:
- 20 Staphylococcal food poisoning, scalded skin syndrome and toxic shock syndrome.
- 25

- Riboflavin (vitamin B2) is a member of the B complex of vitamins which function as coenzymes in metabolic reactions. Riboflavin has two coenzyme forms, flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD) which act in oxidation-reduction reactions such as the cytochrome system of electron transport and the oxidative degradation of pyruvate, fatty acids and amino acids. The first committed step in the biosynthesis of riboflavin is the opening
- 30

of the imidazole ring of GTP. In the presence of 3 H<sub>2</sub>O and Mg<sup>++</sup>, the C-8 of GTP is released as formate accompanied by the release of pyrophosphate by the action of GTP cyclohydrolase II (GCH2; EC 3.5.4.25). This enzyme function is encoded by *ribA* in bacteria and *rib1* in yeast species. Through a series of steps, involving 3,4-dihydroxy-2-butanone 4 phosphate synthase (*ribA*), 6,7-dimethyl-8-ribityllumazine synthase (*ribH*), riboflavin synthase (*ribB*), pyrimidine deaminase and pyrimidine reductase (*ribG*), enzymes encoded by genes within the riboflavin biosynthesis operon, riboflavin is formed. Because the genes required for riboflavin biosynthesis are present in many pathogenic microorganisms, and since riboflavin biosynthesis has shown to be required for virulence in the swine pathogen *Actinobacillus pleuropneumoniae* (Fuller, TE, *et al.* (1996) A riboflavin auxotroph of *Actinobacillus pleuropneumoniae* is attenuated in swine. Infect. Immun. 64:4659-4664), these gene products represent broad spectrum antibacterial as well as antifungal targets.

The frequency of *Staphylococcus aureus* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Staphylococcus aureus* strains which are resistant to some or all of the standard antibiotics. This phenomenon has created a demand for both new anti-microbial agents, vaccines, and diagnostic tests for this organism.

Clearly, there exists a need for factors, such as the ribG embodiments of the invention, that have a present benefit of being useful to screen compounds for antibiotic activity. Such factors are also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their antagonists and agonists to find ways to prevent, ameliorate or correct such infection, dysfunction and disease.

Certain of the polypeptides of the invention possess amino acid sequence homology to a known *Bacillus subtilis* ribG protein.

#### SUMMARY OF THE INVENTION

It is an object of the invention to provide polypeptides that have been identified as novel ribG polypeptides by homology between the amino acid sequence set out in Table 1 [SEQ ID NO:

2 or 4] and a known amino acid sequence or sequences of other proteins such as *Bacillus subtilis* ribG protein.

It is a further object of the invention to provide polynucleotides that encode ribG polypeptides, particularly polynucleotides that encode the polypeptide herein designated ribG.

5 In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding ribG polypeptides comprising a sequence set out in Table 1 [SEQ ID NO:1 or 3] which includes a full length gene, or a variant thereof.

In another particularly preferred embodiment of the invention there is a novel ribG protein from *Staphylococcus aureus* comprising the amino acid sequence of Table 1 [SEQ ID  
10 NO:2 or 4], or a variant thereof.

In accordance with another aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Staphylococcus aureus* WCUH29 on deposit strain, which nucleic acid is contained in the deposited strain.

A further aspect of the invention there are provided isolated nucleic acid molecules  
15 encoding ribG, particularly *Staphylococcus aureus* ribG, including mRNAs, cDNAs, genomic DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

In accordance with another aspect of the invention, there is provided the use of a  
20 polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization. Among the particularly preferred embodiments of the invention are naturally occurring allelic variants of ribG and polypeptides encoded thereby.

Another aspect of the invention there are provided novel polypeptides of *Staphylococcus aureus* referred to herein as ribG as well as biologically, diagnostically, prophylactically,  
25 clinically or therapeutically useful variants thereof, and compositions comprising the same.

Among the particularly preferred embodiments of the invention are variants of ribG polypeptide encoded by naturally occurring alleles of the ribG gene.

In a preferred embodiment of the invention there are provided methods for producing the aforementioned ribG polypeptides.

30 In accordance with yet another aspect of the invention, there are provided inhibitors to such polypeptides, useful as antibacterial agents, including, for example, antibodies.

In accordance with certain preferred embodiments of the invention, there are provided products, compositions and methods for assessing ribG expression, treating disease, assaying genetic variation, and administering a ribG polypeptide or polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Staphylococcus aureus* bacteria.

5 In accordance with certain preferred embodiments of this and other aspects of the invention there are provided polynucleotides that hybridize to ribG polynucleotide sequences, particularly under stringent conditions.

In certain preferred embodiments of the invention there are provided antibodies against ribG polypeptides.

10 In other embodiments of the invention there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity of a polypeptide or polynucleotide of the invention comprising: contacting a polypeptide or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other interaction between the compound and the polypeptide or polynucleotide to  
15 assess the binding to or other interaction with the compound, such binding or interaction being associated with a second component capable of providing a detectable signal in response to the binding or interaction of the polypeptide or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity of the polypeptide or polynucleotide by detecting the presence or absence of a signal generated from  
20 the binding or interaction of the compound with the polypeptide or polynucleotide.

In accordance with yet another aspect of the invention, there are provided ribG agonists and antagonists, preferably bacteriostatic or bacteriocidal agonists and antagonists.

In a further aspect of the invention there are provided compositions comprising a ribG polynucleotide or a ribG polypeptide for administration to a cell or to a multicellular organism.

25 Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

## DESCRIPTION OF THE INVENTION

The invention relates to novel ribG polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a novel ribG of *Staphylococcus aureus*, which is related by amino acid sequence homology to *Bacillus subtilis* ribG polypeptide. The invention relates especially to ribG having the nucleotide and amino acid sequences set out in Table 1 as SEQ ID NO: 1 and SEQ ID NO: 2 respectively ribG.

TABLE 1

ribG Polynucleotide and Polypeptide Sequences

10

(A) Sequences from *Staphylococcus aureus* ribG polynucleotide sequence [SEQ ID NO:1].  
5' -

15

20

25

ATGGATTATGCGATTCAACTTGCAAATATGGTACAAGGTCAAACAGGTGTTAATCCACCCGTTGGCGCTG  
TTGTAGTTAATGAAGGTAGGATTGTTGGTATTGGTGCACACTTGAGAAAAGGTGACAAGCATGCGGAGGT  
TCAAGCACTTGATATGGCACAACAAAATGCTGAAGGTGCGACGATTTATATTACGTTAGAGCCATGTAGT  
CATTTTGGTTCAACACCACCCTGTGTTAACAAAATTATTGATTGTAAGATAGCAAAAGTAGTATACGCAA  
CAAAAGACAATTCGTTAGACACACATGGTGATGAGACGTTACGGGCTCACGGTATTGAGGTTGAATGCGT  
TGATGATGAACGGGCATCACAATTATACCAAGACTTTTTTAAAGCAAAAGCAAAAGCAACTGCCACAAATT  
ACAGTGAAAGTATCTGCAAGTTTAGATGGTAAACAAGCGAATGATAATGGACAAAGTCAATGGATTACTA  
ACAAAGAGGTTAAACAAGATGTCTATAAGTTAAGACATCGACACGACGCAGTGTTAACTGGAAGACGTAC  
AGTTGAATTAGATGATCCACAATATACTACACGTATTCAAGATGGAAAAACCCTATAAAAAGTAATATTG  
TCTAAGTCTGGGAATATTCAATTTAATCAGCAAATTTATCAAGATGAATCAACACCAATTTGGATATATA  
CTGAAATCCAAATTTAACAAGCAATCAAACACATATTGAAATTATTTACTTGAAAGTCTTGTGATTTAAC  
AACAATTCCTCACAATTTATATAAAAGAGGAGTTGGAACCTTGTCTAGTCGAGGCAGGTCCAACCACTACT  
TCAGAATTCTCCATCTATTATATAGATGAATTTATTCTCTATTATGCCCCGAAATTAATTGGCGGATCTG  
GAAATTATCAATTTTATCAAACAAATGATGTGATTGAGATACCAGATGCGAACCAATTTGAAATTGTTCA  
TTCCGAGTTATTAAATCAAATGTAAATTAACCTTTACGAAAGAAGTGA-3'

30

(B) *Staphylococcus aureus* ribG polypeptide sequence deduced from the polynucleotide  
sequence in this table [SEQ ID NO:2].

NH<sub>2</sub>-

MDYAIQLANMVQGQTGVNPPVGAVVVNEGRIVGIGALRLKGDKHAEVQALDMAQQNAEGATIIYITLEPCS  
HFGSTPPCVNKIIDCKIAKVYATKDNSLDTHGDETLRAHGIEVECVDDERASQLYQDFFKAKAKQLPQI  
TVKVSASLDGKQANDNGSQWITNKEVKQDVYKLRHRHDAVLGTGRRTVELDDPQYTTRIQDGKNPIKVIL

SKSGNIHFNQIYQDESTPIWIYTENPNLTSNQTHIEIYLKSCDLTTILHNLTKRGVGTLLVEAGPTTT  
SEFSIYYIDEFILYYAPKLIGSGSNYQFYQTNDVIEIPDANQFEIVHSELLNQNVKLTLRKK-COOH

(C) Polynucleotide sequences comprising *Staphylococcus aureus* ribG ORF sequence  
5 [SEQ ID NO:3].

5' -

AANCACCAATCCNATTGGGAGGNAATCCAAATCAATNCCCGGANNCCCAATCCAAGTTAATTAAGTCCAA  
GGTTTTGGAACATTACCAAATATGATTCCGATGAGGTCAAATGNCAANCGGTGTTAATAAACTACGAAAT  
GNTGTGNAAATGATAGTAGANCAAGTTGCGCATACAGTNTCTCNATTATATGATGCTTTAGAATCGAATG  
10 AGCAACAACAGCGCAGTTACAATCAATAATTTGTAAGTAAAGATAATAAAGAGAACGCTCTATAGAGAC  
GAATTGAAGGTTTGATTTTAATGTCTGTTAGTAAGAATCATATCAATGAGATGCCTATAGTACTCAGATT  
ATATTAAATTAAAACCGTCATTAATTGTTTTTTTAGAAAACATATAGTATCATTTTTAAATGTAGTTGACA  
TACTACGTACTCAAATAATCTATAACAATTTTATATATAATTCTTTTCGGGGCAGGGTGAAATTCCTCAACC  
GGCAGTAAATAAAGCCTGCGACCTGCTAATATGTTTCATATTAGTGGCTGATCTAGTGAGATTCTAGAGC  
15 CGACAGTTAAAGTCTGGATGGGAGAAAGAATGTTAATTATCGACAAAGATAATGTAGCGTATTTGTAAAA  
ATGTGTACAAATAGGCTTATTTAACGATAA  
ATTTTTCTCCTTTGCATCTTAATTCATGATGTGAGGATTTTTTGTATTATAGAGGTGATCATTGAGTCAA  
TTTATGGATTATGCGATTCAACTTCCAAATATGGTACAAGGTCANACAGGTGTTAATCCACCCGTTGGCG  
CTGTTGTAGTTAATGAAGGTAGGATTGTTGGTATTGGTGCACACTTGAGAAAAGGTGACAAGCATGCGGA  
20 GGTTCAGCACTTGATATGGCACAACANAATGCTGAAGGTGCGACGATTTATATTACGTTAGAGCCATGT  
AGTCATTTTGGTTCAACACCACCCTGTGTTAACAAAATTATTGATTGTAAGATAGCANAAGTAGTATTAC  
NCAACANAAGACAATTCCGTTAGACACACATGGGTGATGAGACGTTACGGGGCTCCACGGTATTTGAGGG  
TTGAATTGCGTTGGATGATGAACGGGCATCACAATTATACCAAAGACTTTTTTTAAAGCAAAAAGCAAAG  
CAACTTGCCACAAAATTACAGTGAAAGTNTCTTGAAAGTTTAGATGGGTAAACAAAGCGAATTGATAATG  
25 GACAAAGTCAATGGATTACTAACAAAGAGGTTAAACAAGATGTCTATAG-3'

(D) *Staphylococcus aureus* ribG polypeptide sequence deduced from the polynucleotide  
ORF sequence in this table [SEQ ID NO:4].

NH<sub>2</sub>-

30 MDYAIQLPNMVQGXTVGNPPVGAVVVNEGRIVGIGAHLRKGDKHAQVQALDMAQXNAEGATIIYITLEPCS  
HFGSTPPCVNKKIIDCKIAXVVLXNXRQFR -COOH

---

#### Deposited materials

35 A deposit containing a *Staphylococcus aureus* WCUH 29 strain has been deposited with  
the National Collections of Industrial and Marine Bacteria Ltd. (herein "NCIMB"), 23 St. Machar  
Drive, Aberdeen AB2 1RY, Scotland on 11 September 1995 and assigned NCIMB Deposit No.

40771, and referred to as *Staphylococcus aureus* WCUH29 on deposit. The *Staphylococcus aureus* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

The deposited strain contains the full length ribG gene. The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

The deposit of the deposited strain has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

A license may be required to make, use or sell the deposited strain, and compounds derived therefrom, and no such license is hereby granted.


One aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Staphylococcus aureus* WCUH 29 strain contained in the deposited strain. Further provided by the invention are ribG nucleotide sequences of the DNA in the deposited strain and amino acid sequences encoded thereby. Also provided by the invention are ribG polypeptide sequences isolated from the deposited strain and amino acid sequences derived therefrom.

#### **Polypeptides**

The polypeptides of the invention include a polypeptide of Table 1 [SEQ ID NO:2 or 4] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of ribG, and also those which have at least 70% identity to a polypeptide of Table 1 [SEQ ID NO:1 or 3] or the relevant portion, preferably at least 80% identity to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and more preferably at least 90% similarity (more preferably at least 90% identity) to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and still more preferably at least 95% similarity (still more preferably at least 95% identity) to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and also include portions of such polypeptides with

such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

The invention also includes polypeptides of the formula:



5 wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal,  $R_1$  and  $R_3$  are any amino acid residue, m is an integer between 1 and 1000 or zero, n is an integer between 1 and 1000 or zero, and  $R_2$  is an amino acid sequence of the invention, particularly an amino acid sequence selected from Table 1. In the formula above  $R_2$  is oriented so that its amino terminal residue is at the left, bound to  $R_1$ , and its carboxy terminal residue is at the right, bound to  $R_3$ . Any stretch of amino acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

15 A fragment is a variant polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned polypeptides. As with ribG polypeptides fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region, a single larger polypeptide.

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of Table 1 [SEQ ID NO:2 or 4], or of variants thereof, such as a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus. Degradation forms of the polypeptides of the invention in a host cell, particularly a *Staphylococcus aureus*, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha-amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Also preferred are biologically active fragments which are those fragments that mediate activities of ribG, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments



comprising receptors or domains of enzymes that confer a function essential for viability of *Staphylococcus aureus* or the ability to initiate, or maintain cause disease in an individual, particularly a human.

5 Variants that are fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

10 In addition to the standard single and triple letter representations for amino acids, the term "X" or "Xaa" may also be used in describing certain polypeptides of the invention. "X" and "Xaa" mean that any of the twenty naturally occurring amino acids may appear at such a designated position in the polypeptide sequence.

#### **Polynucleotides**

15 Another aspect of the invention relates to isolated polynucleotides, including the full length gene, that encode the ribG polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2 or 4] and polynucleotides closely related thereto and variants thereof.

20 Using the information provided herein, such as a polynucleotide sequence set out in Table 1 [SEQ ID NO:1 or 3], a polynucleotide of the invention encoding ribG polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Staphylococcus aureus* WCUH 29 cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a sequence given in Table 1 [SEQ ID NO:1 or 3], typically a library of clones of chromosomal DNA of *Staphylococcus aureus* WCUH 29 in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent conditions. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently, such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and 25 Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening 30

By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention, the polynucleotide set out in Table 1 [SEQ ID NO:1 or 3] was discovered in a DNA library derived from *Staphylococcus aureus* WCUH 29.


The DNA sequence set out in Table 1 [SEQ ID NO:1 or 3] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID NO:2 or 4] with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known in the art. The polynucleotide of SEQ ID NO: 1, between nucleotide number 1 and the stop codon which begins at nucleotide number 1027 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

RibG of the invention is structurally related to other proteins of the riboflavin specific deaminase family.

The invention provides a polynucleotide sequence identical over its entire length to a coding sequence in Table 1 [SEQ ID NO:1 or 3]. Also provided by the invention is the coding sequence for the mature polypeptide or a fragment thereof, by itself as well as the coding sequence for the mature polypeptide or a fragment in reading frame with other coding sequence, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence. The polynucleotide may also contain non-coding sequences, including for example, but not limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence which encode additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA tag (Wilson *et al.*, *Cell* 37: 767 (1984). Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

A preferred embodiment of the invention is a polynucleotide of comprising nucleotide 1 to the nucleotide immediately upstream of or including nucleotide 1027 set forth in SEQ ID NO:1 of Table 1, both of which encode the ribG polypeptide.

The invention also includes polynucleotides of the formula:



wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal,  $R_1$  and  $R_3$  is any nucleic acid residue, m is an integer between 1 and 3000 or zero, n is an integer between 1 and 3000 or zero, and  $R_2$  is a nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from Table 1. In the polynucleotide formula above  $R_2$  is oriented so that its 5' end residue is at the left, bound to  $R_1$ , and its 3' end residue is at the right, bound to  $R_3$ . Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer. In a preferred embodiment m and/or n is an integer between 1 and 1000.

It is most preferred that the polynucleotides of the inventions are derived from *Staphylococcus aureus*, however, they may preferably be obtained from organisms of the same taxonomic genus. They may also be obtained, for example, from organisms of the same taxonomic family or order.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Staphylococcus aureus* ribG having an amino acid sequence set out in Table 1 [SEQ ID NO:2 or 4]. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or an insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode for variants of the polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2 or 4]. Variants that are fragments of the polynucleotides of the invention may be used to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding ribG variants, that have the amino acid sequence of ribG polypeptide of Table 1 [SEQ ID NO:2 or 4] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of ribG.

Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding ribG polypeptide having an amino acid sequence set out in Table 1 [SEQ ID NO:2 or 4], and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding ribG polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of Table 1 [SEQ ID NO:1 or 3].

The invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or

a fragment thereof; and isolating said DNA sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers described elsewhere herein.

As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding ribG and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to the ribG gene. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

For example, the coding region of the ribG gene may be isolated by screening using a DNA sequence provided in Table 1 [SEQ ID NO: 1 or 3] to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays.

Polynucleotides of the invention that are oligonucleotides derived from the sequences of Table 1 [SEQ ID NOS:1 or 2 or 3 or 4] may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that may encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

5 In addition to the standard A, G, C, T/U representations for nucleic acid bases, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA bases may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a base that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of  
10 generating a premature termination codon in such reading frame.

In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more  
15 prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

#### **Vectors, host cells, expression**

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the  
20 production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a  
25 polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and  
30 infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, enterococci *E. coli*, streptomyces and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and  
5 plant cells.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, *e.g.*, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as  
10 baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express  
15 polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into  
20 the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid  
25 extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during  
30 isolation and or purification.

#### Diagnostic Assays

This invention is also related to the use of the ribG polynucleotides of the invention for use as diagnostic reagents. Detection of ribG in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans, particularly those  
5 infected or suspected to be infected with an organism comprising the ribG gene may be detected at the nucleic acid level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification technique prior  
10 to analysis. RNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of prokaryote present in an individual, may be made by an analysis of the genotype of the prokaryote gene. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence. Point mutations can be identified by hybridizing amplified DNA to labeled  
15 ribG polynucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in the electrophoretic mobility of the DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be  
20 revealed by nuclease protection assays, such as RNase and S1 protection or a chemical cleavage method. See, e.g., Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

Cells carrying mutations or polymorphisms in the gene of the invention may also be detected at the DNA level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations. It is particularly preferred to used RT-PCR in  
25 conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to a nucleic acid encoding ribG can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 2.

30

Table 2

**Primers for amplification of ribG polynucleotides**




SEQ ID NOPRIMER SEQUENCE

5                    5' -ATGAAGGTAGGATTGTTGGTA-3'  
6                    5' -AGTCTTGGTATAATTGTGATGC-3'

5

The invention also includes primers of the formula:



wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R<sub>1</sub> and R<sub>3</sub> is any nucleic acid residue, m is an integer between 1 and 20 or zero, n is an integer between 1 and 20 or zero, and R<sub>2</sub> is a primer sequence of the invention, particularly a primer sequence selected from Table 2. In the polynucleotide formula above R<sub>2</sub> is oriented so that its 5' end residue is at the left, bound to R<sub>1</sub>, and its 3' end residue is at the right, bound to R<sub>3</sub>. Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer being complementary to a region of a polynucleotide of Table 1. In a preferred embodiment m and/or n is an integer between 1 and 10.

The invention further provides these primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying ribG DNA isolated from a sample derived from an individual. The primers may be used to amplify the gene isolated from an infected individual such that the gene may then be subject to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be detected and used to diagnose infection and to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections by *Staphylococcus aureus*, comprising determining from a sample derived from an individual a increased level of expression of polynucleotide having a sequence of Table 1 [SEQ ID NO: 1 or 3]. Increased or decreased expression of ribG polynucleotide can be measured using any on of the methods well known in the art for the quantation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of ribG protein compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a ribG protein, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

### **Antibodies**

The polypeptides of the invention or variants thereof, or cells expressing them can be used as an immunogen to produce antibodies immunospecific for such polypeptides. "Antibodies" as used herein includes monoclonal and polyclonal antibodies, chimeric, single chain, simianized antibodies and humanized antibodies, as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

Antibodies generated against the polypeptides of the invention can be obtained by administering the polypeptides or epitope-bearing fragments, analogues or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies.

Alternatively phage display technology may be utilized to select antibody genes with binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-ribG or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. *et al.*, (1991) *Nature* 352, 624-628).

If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides to purify the polypeptides by affinity chromatography.

Thus, among others, antibodies against ribG- polypeptide may be employed to treat infections, particularly bacterial infections.

5 Polypeptide variants include antigenically, epitopically or immunologically equivalent variants that form a particular aspect of this invention. The term "antigenically equivalent derivative" as used herein encompasses a polypeptide or its equivalent which will be specifically recognized by certain antibodies which, when raised to the protein or polypeptide according to the invention, interfere with the immediate physical interaction between pathogen  
10 and mammalian host. The term "immunologically equivalent derivative" as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

The polypeptide, such as an antigenically or immunologically equivalent derivative or  
15 a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically  
20 or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized"; where the complementarity determining region(s) of the hybridoma-derived  
25 antibody has been transplanted into a human monoclonal antibody, for example as described in Jones, P. et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles  
30 (Wolff et al., *Hum Mol Genet* 1992, 1:363, Manthorpe et al., *Hum. Gene Ther.* 1993:4, 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem.* 1989:

264,16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, 1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* 1989:243,375), particle bombardment (Tang et al., *Nature* 1992, 356:152, Eisenbraun et al., *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS USA* 1984:81,5849).

#### **Antagonists and agonists - assays and molecules**

Polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, e.g., Coligan et al., *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of ribG polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bacteriocidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising ribG polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a ribG agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the ribG polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of ribG polypeptide are most likely to be good antagonists. Molecules that bind well and increase the rate of product production from substrate are agonists. Detection of the rate or level of production of product from substrate may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in ribG polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for ribG antagonists is a competitive assay that combines ribG and a potential antagonist with ribG-binding molecules, recombinant ribG binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. RibG can be labeled, such as by radioactivity or a

colorimetric compound, such that the number of ribG molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing ribG-induced activities, thereby preventing the action of ribG by excluding ribG from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of ribG.

Each of the DNA sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide or inhibitor of the invention to interfere with the initial physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block ribG protein-mediated mammalian cell invasion by, for example, initiating phosphorylation of mammalian tyrosine kinases (Rosenshine *et al.*, *Infect. Immun.* 60:2211 (1992); to block bacterial adhesion between mammalian extracellular matrix

proteins and bacterial ribG proteins that mediate tissue damage and; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

The antagonists and agonists of the invention may be employed, for instance, to inhibit and treat diseases.

*Helicobacter pylori* (herein *H. pylori*) bacteria infect the stomachs of over one-third of the world's population causing stomach cancer, ulcers, and gastritis (International Agency for Research on Cancer (1994) Schistosomes, Liver Flukes and *Helicobacter Pylori* (International Agency for Research on Cancer, Lyon, France; <http://www.uicc.ch/ecp/ecp2904.htm>). Moreover, the international Agency for Research on Cancer recently recognized a cause-and-effect relationship between *H. pylori* and gastric adenocarcinoma, classifying the bacterium as a Group I (definite) carcinogen. Preferred antimicrobial compounds of the invention (agonists and antagonists of ribG) found using screens provided by the invention, particularly broad-spectrum antibiotics, should be useful in the treatment of *H. pylori* infection. Such treatment should decrease the advent of *H. pylori*-induced cancers, such as gastrointestinal carcinoma. Such treatment should also cure gastric ulcers and gastritis.

#### Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with ribG, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Staphylococcus aureus* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector to direct expression of ribG, or a fragment or a variant thereof, for expressing ribG, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual from disease, whether that disease is already established within the individual or not. One way of administering the gene is by accelerating it into the desired cells as a coating on

particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid.

5 A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response, induces an immunological response in such individual to a ribG or protein coded therefrom, wherein the composition comprises a recombinant ribG or protein coded therefrom comprising DNA which codes for and expresses an antigen of said ribG or protein coded therefrom. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T  
10 cells.

A ribG polypeptide or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from  
15 *Hemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

20 Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or  
25 particular fragments thereof which have been shown to encode non-variable regions of bacterial cell surface proteins in DNA constructs used in such genetic immunization experiments in animal models of infection with *Staphylococcus aureus* will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of  
30 monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic

treatments of bacterial infection, particularly *Staphylococcus aureus* infection, in mammals, particularly humans.

5 The polypeptide may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or connective tissue caused, e.g., by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

10 The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the  
15 formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may  
20 also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

25 While the invention has been described with reference to certain ribG protein, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

#### **Compositions, kits and administration**

30 The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or their agonists or antagonists. The polypeptides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject.



Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

Alternatively the composition may be formulated for topical application for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e., devices that are introduced to the body of an individual and remain in position for an extended time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory  
5 peritoneal dialysis (CAPD) catheters.

The composition of the invention may be administered by injection to achieve a systemic effect against relevant bacteria shortly before insertion of an in-dwelling device. Treatment may be continued after surgery during the in-body time of the device. In addition, the composition could also be used to broaden perioperative cover for any surgical technique to  
10 prevent bacterial wound infections, especially *Staphylococcus aureus* wound infections.

Many orthopaedic surgeons consider that humans with prosthetic joints should be considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia. Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to  
15 extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis.

20 Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5  
25 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Each reference disclosed herein is incorporated by reference herein in its entirety. Any  
30 patent application to which this application claims priority is also incorporated by reference herein in its entirety.

## GLOSSARY

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Disease(s)" means and disease caused by or related to infection by a bacteria, including  
5 disease, such as, infections of the upper respiratory tract (e.g., otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g., empyema, lung abscess), cardiac (e.g., infective endocarditis), gastrointestinal (e.g., secretory diarrhoea, splenic absces, retroperitoneal abscess), CNS (e.g., cerebral abscess), eye (e.g., blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis), kidney and urinary tract (e.g.,  
10 epididymitis, intrarenal and perinephric absces, toxic shock syndrome), skin (e.g., impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis) bone and joint (e.g., septic arthritis, osteomyelitis).

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

15 "Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods,  
20 including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis*  
25 *Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., SIAM *J. Applied Math.*, 48: 1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two  
30 sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Atschul, S.F. et

al., *J. Molec. Biol.* 215: 403-410 (1990). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., *et al.*, NCBI NLM NIH Bethesda, MD 20894; Altschul, S., *et al.*, *J. Mol. Biol.* 215: 403-410 (1990). As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having an amino acid sequence having at least, for example, 95% identity to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is

"isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

5 "Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules  
10 comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically  
15 involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or  
20 modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and  
25 RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains  
30 generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such

as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

"Variant(s)" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide

changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

### EXAMPLES

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

#### Example 1 Strain selection, Library Production and Sequencing

The polynucleotide having a DNA sequence given in Table 1 [SEQ ID NO:1 or 3] was obtained from a library of clones of chromosomal DNA of *Staphylococcus aureus* in *E. coli*. The sequencing data from two or more clones containing overlapping *Staphylococcus aureus* DNAs was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example:

Methods 1 and 2 below.

Total cellular DNA is isolated from *Staphylococcus aureus* WCUH 29 according to standard procedures and size-fractionated by either of two methods.

#### Method 1

Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

## Method 2

Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., RsaI, PstI, AluI, BshI235I), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

### Example 2 ribG Characterization

#### Riboflavin Biosynthesis Operon

This ORF is part of an operon which encodes genes *ribG*, *ribB*, *ribA* and *ribH*. Gene *ribG* starts at nucleotide 1 and ends at nucleotide 1029. Gene *ribB* starts at nucleotide 1036 and ends at nucleotide 1668. Gene *ribA* starts at nucleotide 1679 and ends at nucleotide 2860. Gene *ribH* starts at nucleotide 2873 and ends at nucleotide 3337. The operon sequence below is SEQ ID NO:7.

ATGGATTATGCGATTCAACTTGCAAATATGGTACAAGGTCAAACAGGTGTTAATCCACCCGTTGGCGCTG  
TTGTAGTTAATGAAGGTAGGATTGTTGGTATTGGTGCACACTTGAGAAAAGGTGACAAGCATGCGGAGGT  
TCAAGCACTTGATATGGCACAACAAATGCTGAAGGTGCGACGATTTATATTACGTTAGAGCCATGTAGT  
CATTTTGGTTCAACACCACCCTGTGTTAACAAAATTATTGATTGTAAGATAGCAAAAGTAGTATACGCAA  
CAAAAGACAATTCGTTAGACACACATGGTGATGAGACGTTACGGGCTCACGGTATTGAGGTTGAATGCGT  
TGATGATGAACGGGCATCACAATTATACCAAGACTTTTTTAAAGCAAAAGCAAAGCAACTGCCACAAATT  
ACAGTGAAAGTATCTGCAAGTTTAGATGGTAAACAAGCGAATGATAATGGACAAAGTCAATGGATTACTA  
ACAAAGAGGTTAAACAAGATGTCTATAAGTTAAGACATCGACACGACGCAGTGTTAACTGGAAGACGTAC  
AGTTGAATTAGATGATCCACAATATACTACACGTATTCAAGATGGAAAAACCCTATAAAAGTAATATTG  
TCTAAGTCTGGGAATATTCAATTTAATCAGCAAATTTATCAAGATGAATCAACACCAATTTGGATATATA  
CTGAAAATCCAAATTTAACAAGCAATCAAACACATATTGAAATTATTTACTTGAAGTCTTGTGATTTAAC  
AACAATTCTTCACAATTTATATAAAAGAGGAGTTGGAACTTTGCTAGTCGAGGCAGGTCCAACCACTACT  
TCAGAATTCTCCATCTATTATATAGATGAATTTATTCTCTATTATGCCCCGAAATTAATTGGCGGATCTG  
GAAATTATCAATTTTATCAAACAAATGATGTGATTGAGATACCAGATGCGAACCAATTTGAAATTGTTCA  
TTCCGAGTTATTAAATCAAAATGTTAAATTAACTTTACGAAAGAAGTGATGATGCATGTTTACTGGCATC  
GTTGAAGAAATAGGTGTCGTTAAAAGTGTTCAAATTCGTCAATCAGTCAGGACGATTGAAATCGAAGCAC





[illegible]

(i) APPLICANT: Palmer, Leslie M.

(iii) NUMBER OF SEQUENCES: 7

(A) ADDRESSEE: Dechert Price & Rhoads

(B) STREET: 4000 Bell Atlantic Tower, 1717 Arch Stre

(C) CITY: Philadelphia

(D) STATE: PA

(E) COUNTRY: US

(F) ZIP: 19103

(A) MEDIUM TYPE: Diskette

(B) COMPUTER: IBM Compatible

(C) OPERATING SYSTEM: DOS

(D) SOFTWARE: FastSEQ for Windows Version 2.0

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(A) NAME: Dickinson, Todd O

(B) REGISTRATION NUMBER: 28,354  
(C) REFERENCE/DOCKET NUMBER: P50444-9

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 215-994-2252  
(B) TELEFAX: 215-994-2222  
(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1029 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGGATTATG CGATTCAACT TGCAAATATG GTACAAGGTC AAACAGGTGT TAATCCACCC 60  
GTTGGCGCTG TTGTAGTTAA TGAAGGTAGG ATTGTTGGTA TTGGTGCACA CTTGAGAAAA 120  
GGTGACAAGC ATGCGGAGGT TCAAGCACTT GATATGGCAC AACAAAATGC TGAAGGTGCG 180  
ACGATTTATA TTACGTTAGA GCCATGTAGT CATTTTGGTT CAACACCACC CTGTGTTAAC 240  
AAAATTATTG ATTGTAAGAT AGCAAAAGTA GTATACGCAA CAAAAGACAA TTCGTTAGAC 300  
ACACATGGTG ATGAGACGTT ACGGGCTCAC GGTATTGAGG TTGAATGCGT TGATGATGAA 360  
CGGGCATCAC AATTATACCA AGACTTTTTT AAAGCAAAAG CAAAGCAACT GCCACAAATT 420  
ACAGTGAAAAG TATCTGCAAG TTTAGATGGT AAACAAGCGA ATGATAATGG ACAAAGTCAA 480  
TGGATTACTA ACAAAGAGGT TAAACAAGAT GTCTATAAGT TAAGACATCG ACACGACGCA 540  
GTGTTAACTG GAAGACGTAC AGTTGAATTA GATGATCCAC AATATACTAC ACGTATTCAA 600  
GATGGAAAAA ACCCTATAAA AGTAATATTG TCTAAGCTG GGAATATTCA TTTTAATCAG 660  
CAAATTTATC AAGATGAATC AACACCAATT TGGATATATA CTGAAAATCC AAATTTAACA 720  
AGCAATCAAA CACATATTGA AATTATTTAC TTGAAGTCTT GTGATTTAAC AACAATTCTT 780  
CACAATTTAT ATAAAAGAGG AGTTGGAAC T TGCTAGTCG AGGCAGGTCC AACCCTACT 840  
TCAGAATTCT CCATCTATTA TATAGATGAA TTTATTCTCT ATTATGCCCC GAAATTAATT 900  
GGCGGATCTG GAAATTATCA ATTTTATCAA ACAAATGATG TGATTGAGAT ACCAGATGCG 960  
AACCAATTTG AAATTGTTCA TTCCGAGTTA TTAAATCAAA ATGTTAAATT AACTTTACGA 1020  
AAGAAGTGA 1029

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 342 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Asp Tyr Ala Ile Gln Leu Ala Asn Met Val Gln Gly Gln Thr Gly  
1 5 10 15  
Val Asn Pro Pro Val Gly Ala Val Val Val Asn Glu Gly Arg Ile Val  
20 25 30  
Gly Ile Gly Ala His Leu Arg Lys Gly Asp Lys His Ala Glu Val Gln  
35 40 45  
Ala Leu Asp Met Ala Gln Gln Asn Ala Glu Gly Ala Thr Ile Tyr Ile  
50 55 60  
Thr Leu Glu Pro Cys Ser His Phe Gly Ser Thr Pro Pro Cys Val Asn  
65 70 75 80  
Lys Ile Ile Asp Cys Lys Ile Ala Lys Val Val Tyr Ala Thr Lys Asp  
85 90 95  
Asn Ser Leu Asp Thr His Gly Asp Glu Thr Leu Arg Ala His Gly Ile  
100 105 110  
Glu Val Glu Cys Val Asp Asp Glu Arg Ala Ser Gln Leu Tyr Gln Asp  
115 120 125  
Phe Phe Lys Ala Lys Ala Lys Gln Leu Pro Gln Ile Thr Val Lys Val  
130 135 140  
Ser Ala Ser Leu Asp Gly Lys Gln Ala Asn Asp Asn Gly Gln Ser Gln  
145 150 155 160  
Trp Ile Thr Asn Lys Glu Val Lys Gln Asp Val Tyr Lys Leu Arg His  
165 170 175  
Arg His Asp Ala Val Leu Thr Gly Arg Arg Thr Val Glu Leu Asp Asp  
180 185 190  
Pro Gln Tyr Thr Thr Arg Ile Gln Asp Gly Lys Asn Pro Ile Lys Val  
195 200 205  
Ile Leu Ser Lys Ser Gly Asn Ile His Phe Asn Gln Gln Ile Tyr Gln  
210 215 220  
Asp Glu Ser Thr Pro Ile Trp Ile Tyr Thr Glu Asn Pro Asn Leu Thr  
225 230 235 240

Ser	Asn	Gln	Thr	His	Ile	Glu	Ile	Ile	Tyr	Leu	Lys	Ser	Cys	Asp	Leu	
				245				250				255				
Thr	Thr	Ile	Leu	His	Asn	Leu	Tyr	Lys	Arg	Gly	Val	Gly	Thr	Leu	Leu	
				260				265				270				
Val	Glu	Ala	Gly	Pro	Thr	Thr	Thr	Ser	Glu	Phe	Ser	Ile	Tyr	Tyr	Ile	
				275				280				285				
Asp	Glu	Phe	Ile	Leu	Tyr	Tyr	Ala	Pro	Lys	Leu	Ile	Gly	Gly	Ser	Gly	
				290				295				300				
Asn	Tyr	Gln	Phe	Tyr	Gln	Thr	Asn	Asp	Val	Ile	Glu	Ile	Pro	Asp	Ala	
305				310				315				320				
Asn	Gln	Phe	Glu	Ile	Val	His	Ser	Glu	Leu	Leu	Asn	Gln	Asn	Val	Lys	
				325				330				335				
Leu	Thr	Leu	Arg	Lys	Lys											
				340												

- (2) INFORMATION FOR SEQ ID NO:3:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AANACCAAT	CCNATTGGGA	GGNAATCCAA	ATCAATNCCC	GGANNCCCAA	TCCAAGTTAA	60
TTAAGTCCAA	GGTTTTGGAA	CATTACCAA	TATGATTCCG	ATGAGGTCAA	ATGNCAANCG	120
GTGTTAATAA	ACTACGAAAT	GNTGTGNAAA	TGATAGTAGA	NCAAGTTGCG	CATACAGTNT	180
CTCNATTATA	TGATGCTTTA	GAATCGAATG	AGCAACAACA	GCGCAGTTAC	AATCAATAAT	240
TTGTAAC TAG	AAGATAATAA	AGAGAACGCT	CTATAGAGAC	GAATTGAAGG	TTTGATTTTA	300
ATGTCTGTTA	GTAAGAATCA	TATCAATGAG	ATGCCTATAG	TACTCAGATT	ATATTAAATT	360
AAAACCGTCA	TTAATTGTTT	TTTTAGAAAA	CATATAGTAT	CATTTTAAAT	GTAGTTGACA	420
TACTACGTAC	TCAAATAATC	TATAACAATT	TCATATATAA	TTCTTTCGGG	GCAGGGTGAA	480
ATCCCCAACC	GGCAGTAAAT	AAAGCCTGCG	ACCTGCTAAT	ATGTTTCATA	TTAGTGGCTG	540
ATCTAGTGAG	ATTCTAGAGC	CGACAGTTAA	AGTCTGGATG	GGAGAAAGAA	TGTTAATTAT	600
CGACAAAGAT	AATGTAGCGT	ATTTGTAAAA	ATGTGTACAA	ATAGGCTTAT	TTAACGATAA	660
ATTTTTCTCC	TTTGCATCTT	AATTCATGAT	GTGAGGATTT	TTTGTTTATA	GAGGTGATCA	720
TTTGAGTCAA	TTTATGGATT	ATGCGATTCA	ACTTCCAAAT	ATGGTACAAG	GTCANACAGG	780
TGTTAATCCA	CCCGTTGGCG	CTGTTGTAGT	TAATGAAGGT	AGGATTGTTG	GTATTGGTGC	840

ACACTTGAGA	AAAGGTGACA	AGCATGCGGA	GGTTCAAGCA	CTTGATATGG	CACAACANAA	900
TGCTGAAGGT	GCGACGATTT	ATATTACGTT	AGAGCCATGT	AGTCATTTTG	GTTCAACACC	960
ACCCTGTGTT	AACAAAATTA	TTGATTGTAA	GATAGCANAA	GTAGTATTAC	NCAACANAAG	1020
ACAATTCCGT	TAGACACACA	TGGGTGATGA	GACGTTACGG	GGCTCCACGG	TATTTGAGGG	1080
TTGAATTGCG	TTGGATGATG	AACGGGCATC	ACAATTATAC	CAAAGACTTT	TTTTAAAGCA	1140
AAAAGCAAAG	CAACTTGCCA	CAAAATTACA	GTGAAAGTNT	CTTGAAAGTT	TAGATGGGTA	1200
AACAAAGCGA	ATTGATAATG	GACAAAGTCA	ATGGATTACT	AACAAAGAGG	TTAAACAAGA	1260
TGTCTATAG						1269

(2) INFORMATION FOR SEO ID NO:4:

[illegible]

18	19	20	21	22	23	24	25	26	27	28	29	30
18	19	20	21	22	23	24	25	26	27	28	29	30

[illegible][illegible]

18	19	20	21	22	23	24	25	26	27	28	29	30
18	19	20	21	22	23	24	25	26	27	28	29	30

[illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible]

18	19	20	21	22	23	24	25	26	27	28	29	30
18	19	20	21	22	23	24	25	26	27	28	29	30

[illegible][illegible][illegible][illegible][illegible][illegible]

TGGATTACTA	ACAAAGAGGT	TAAACAAGAT	GTCTATAAGT	TAAGACATCG	ACACGACGCA	540
GTGTTAAGTG	GAAGACGTAC	AGTTGAATTA	GATGATCCAC	AATATACTAC	ACGTATTCAA	600
GATGGAAAAA	ACCCTATAAA	AGTAATATTG	TCTAAGTCTG	GGAATATTCA	TTTTAATCAG	660
CAAATTTATC	AAGATGAATC	AACACCAATT	TGGATATATA	CTGAAAAATCC	AAATTTAACA	720
AGCAATCAAA	CACATATTGA	AATTATTTAC	TTGAAGTCTT	GTGATTTAAC	AACAATTCTT	780
CACAATTTAT	ATAAAAGAGG	AGTTGGAAC	TTGCTAGTCG	AGGCAGGTCC	AACCACTACT	840
TCAGAATTCT	CCATCTATTA	TATAGATGAA	TTTATTCTCT	ATTATGCCCC	GAAATTAATT	900
GGCGGATCTG	GAAATTATCA	ATTTTATCAA	ACAAATGATG	TGATTGAGAT	ACCAGATGCG	960
AACCAATTTG	AAATTGTTCA	TTCCGAGTTA	TTAAATCAAA	ATGTTAAATT	AACCTTACGA	1020
AAGAAGTGAT	GATGCATGTT	TACTGGCATC	GTTGAAGAAA	TAGGTGTCGT	TAAAAGTGTT	1080
CAAATTCGTC	AATCAGTCAG	GACGATTGAA	ATCGAAGCAC	ATAAGATTAC	GGCAGATATG	1140
CATATTGGTG	ATTCCATCAG	TGTTAATGGT	GCATGTTTAA	CAGTGATTGA	TTTTGATCAG	1200
ACATCTTTTA	CTGTTCAGT	AATTAAAAGC	ACTGAAAATA	AAACCTATTT	AGCAGATGTT	1260
AAGCGACAAT	CAGAAGTAAA	TTTAGAGCGT	GCCATGAGTG	GTAACGGTAG	GTTTGGTGGA	1320
CATTTTGTCC	TCGGTCATGT	AGATGAACTA	GGAACAGTTT	CAAAAATAAA	TGAAACAGCC	1380
AATGCCAAAA	TTATTACGAT	TCAATGTAGC	CAACATATTA	ATAATCAGTT	AGTTAAGCAA	1440
GGGTCTATTA	CTGTGGATGG	TGTAAGTCTA	ACGGTATTTG	ATAAGCATGA	TAACAGTTTT	1500
GACATTCATC	TTATTCCAGA	AACGAGGCGT	TCAACGATTT	TATCATCCAA	AAAATTAGGA	1560
GATAAAGTAC	ATTTAGAAAC	AGACGTTTTG	TTTAAATATG	TTGAAAATAT	TTTAAATAAA	1620
GATAAAGACC	AATTATCTGT	AGATAAATTA	AGAGCATTTG	GTTTTTAGGA	GGGGTAGCAT	1680
GCAATTTCGAT	AATATTGACA	GTGCTTTAAT	GGCTTTAAAA	AATGGAGAAA	CAATTATTGT	1740
AGTAGATGAT	GAGAATCGTG	AAAATGAAGG	TGATTTAGTA	GCGGTTACTG	AATGGATGAA	1800
CGATAATACC	ATTAATTTTA	TGGCGAAAGA	AGCAAGGGGA	TTAATATGCG	CACCAGTGTC	1860
TAAAGATATT	GCACAACGTT	TGGATTTGGT	ACAAATGGTT	GATGATAACT	CCGACATCTT	1920
TGGTACGCAA	TTTACAGTGA	GTATTGATCA	TGTAGATACA	ACAACAGGAA	TTAGTGCTTA	1980
TGAACGTACA	TTGACTGCCA	AAAAGCTCAT	TGATCCTAGT	AGTGAAGCTA	AAGATTTTAA	2040
TCGTCCGGT	CATTTATTTT	CATTAGTAGC	ACAAGATAAA	GGCGTATTAG	CTAGAAATGG	2100
ACACACAGAA	GCGGCTGTTG	ATTTAGCTAA	ACTTACTGGT	GCCAAGCCCG	CTGGTGTCAT	2160
TTGTGAGATT	ATGAATGATG	ACGGCACGAT	GGCGAAAGGA	CAAGATTTAC	AAAATTTTAA	2220
AGAAAAACAT	CAATTAAAGA	TGATTACGAT	TGATGATTTA	ATTGAATATC	GTAAAAAATT	2280
AGAACCAGAA	ATTGAATTTA	AGGCAAAAGT	GAAAATGCCT	ACAGATTTCT	GAACATTTGA	2340
TATGTATGGT	TTTAAAGCGA	CATACACAGA	TGAAGAGATA	GTTGTACTGA	CAAAGGTGTC	2400
AATTCGACAA	CATGAAAATG	TACGCTTACA	TTCTGCGTGC	CTTACAGGCG	ATATTTTCCA	2460
TAGTCAACGT	TGTGATTGTG	GTGCTCAACT	TGAATCGTCT	ATGAAGTATA	TCAATGAACA	2520
TGGTGGCATG	ATTATTTATC	TACCTCAAGA	AGGTCGTGGC	ATAGGATTGT	TAAACAAATT	2580
ACGCGCATAT	GAATTAATTG	AGCAAGGATA	TGATACAGTA	ACTGCAAATT	TAGCATTAGG	2640
TTTTGATGAA	GATTTACGAG	ATTATCATAT	TGCTGCACAG	ATTTTAAAT	ATTTTAACAT	2700
CGAACATATC	AATTTATTAA	GTAATAATCC	AAGTAAATTT	GAGGGATTAA	AACAATATGG	2760
CATTGATATT	GCAGAAAGAA	TTGAAGTTAT	CGTACCAGAA	ACGGTACATA	ATCATGATTA	2820
TATGGTAACG	AAAAAAATAA	AAATGGGTCA	TTTAATATAG	GAGGACTTTA	ACATGAATTT	2880



TGAAGGTAAA	TTAATTGGAA	AAGATTTGAA	AGTTGCAATC	GTAGTTAGTC	GATTTAATGA	2940
TTTTATCACT	GGAAGATTAC	TTGAAGGTGC	AAAAGATACT	TTGATTGAC	ATGATGTTAA	3000
TGAAGACAAT	ATTGATGTAG	CATTTGTTCC	TGGTGCGTTT	GAAATTCCTT	TAGTAGCTAA	3060
AAAATTAGCC	TCATCAGGAA	ATTATGATGC	AATAATTACA	TTAGGATGCG	TAATTCGCGG	3120
TGCTACGTCT	CATTATGATT	ATGTTTGTAA	TGAAGTGCGA	AAGGTGTTTC	TAAAGTAAAT	3180
GATCAAATA	ATGTACCAGT	CATATTTGGT	ATTTTAACGA	CTGAAAGTAT	TGAACAAGCT	3240
GTGGAAAGAG	CAGGTACGAA	AGCTGGTAAT	AAAGGTGCCG	AAGCAGCAGT	AAGTGCAATT	3300
GAAATGGCTA	ATTTATTAAA	ATCTATAAAA	GCATAG			3336

668780-449266

**THIS PAGE BLANK (USPTO)**